

# The CSIRO Mk3L climate system model v1.1: PMIP2 experiments

Steven J. Phipps  
CSIRO Marine and Atmospheric Research, Australia  
[steven.phipps@csiro.au](mailto:steven.phipps@csiro.au)

11 July 2008

## 1 Model documentation

### 1.1 Standard reference

The standard reference for the CSIRO Mk3L climate system model is *Phipps (2006)*. A PDF of this report can be downloaded using the following URL:

<http://staff.acecrc.org.au/~sjhipps/publications/phipps2006a.pdf>

### 1.2 Short description

The CSIRO Mk3L climate system model v1.1 is a computationally-efficient coupled atmosphere-sea ice-ocean general circulation model, suitable for studying climate variability and change on millennial timescales. The atmospheric component of Mk3L comprises a spectral general circulation model, a dynamic-thermodynamic sea ice model, and a land surface model with fixed vegetation and soil types. A coarse horizontal resolution of R21 is employed throughout, giving zonal and meridional resolutions of  $5.625^\circ$  and  $\sim 3.18^\circ$  respectively. A hybrid vertical coordinate is used in the atmosphere, with 18 vertical levels. The oceanic component of Mk3L is a coarse-resolution,  $z$ -coordinate general circulation model. There are 21 vertical levels, and the horizontal resolution is double that of the atmospheric component. Each atmosphere model grid cell is exactly matched by four ocean model grid cells, giving zonal and meridional resolutions of  $2.8125^\circ$  and  $\sim 1.59^\circ$  respectively.

### 1.3 Comparison with IPCC versions

The atmospheric, land surface and sea ice components of Mk3L are the same as those of the CSIRO Mk3 climate system model (*Gordon et al., 2002*), which was used to produce simulations for AR4. However, the horizontal resolution is reduced from T63 to R21.

The oceanic component of Mk3L is the same as that of the CSIRO Mk2 coupled model (*Gordon and O'Farrell, 1997; Hirst et al., 2000*), which was used to produce simulations for AR3. However, the horizontal resolution is doubled from  $5.625^\circ \times \sim 3.18^\circ$  to  $2.8125^\circ \times \sim 1.59^\circ$ .

## 1.4 Comparison with v1.0

Relative to the CSIRO Mk3L climate system model v1.0, the horizontal resolution of the oceanic component has been doubled from  $5.625^\circ \times \sim 3.18^\circ$  to  $2.8125^\circ \times \sim 1.59^\circ$ . A new bathymetry has been derived, and the positions of the coastlines have been moved so that they match exactly with the positions of the coastlines on the atmosphere model grid. Many new straits have therefore been opened up. The ocean model has also been re-tuned, with reduced horizontal viscosity and reduced diffusivities.

The experimental design for the PMIP2 experiments differs from that used with v1.0 as follows:

- For the ocean model spin-up run, the convergence criterion for the rate of change in global-mean temperature on each model level has been reduced from  $0.005^\circ\text{C}/\text{century}$  to  $0.001^\circ\text{C}/\text{century}$ .
- For the atmosphere model spin-up run, the sea surface temperatures are derived from the NOAA Optimum Interpolation v2 sea surface temperature analysis (*Reynolds et al., 2002*), rather than the World Ocean Atlas 1998 (*National Oceanographic Data Center, 2002*).
- The mid-Holocene experiment uses an atmospheric  $\text{CO}_2$  concentration of 280 ppm, rather than 277 ppm.

## 1.5 Access to the model source code

The CSIRO Mk3L climate system model is freely available to the research community. Access to the source code can be obtained by completing the form at the following URL:

<http://www.tpac.org.au/main/csiromk3l>

# 2 Description of experiments

## 2.1 Spin-up procedure

### 2.1.1 Ocean model

The World Ocean Atlas 1998 (*National Oceanographic Data Center, 2002*) temperatures and salinities were used to initialise the model, with the velocities being set to zero. The model was forced with climatological NCEP-DOE Reanalysis 2 (*Kanamitsu et al., 2002*) wind stresses, while the temperature and

salinity of the upper layer were relaxed towards the World Ocean Atlas 1998 values using a relaxation timescale of 20 days.

The model was initially integrated to equilibrium using asynchronous timestepping, with a tracer timestep of 1 day and a momentum timestep of 20 minutes. Equilibrium was attained after a total of 6,500 years of asynchronous timestepping, with the convergence criterion being that the rates of change in global-mean temperature and salinity on each model level must be less than  $0.001^{\circ}\text{C}/\text{century}$  and  $0.001 \text{ psu}/\text{century}$  respectively.

The model was then integrated to equilibrium under synchronous timestepping, with a timestep of 1 hour being used to integrate both the tracer and momentum equations. The convergence criterion was satisfied after 500 years.

For the purposes of deriving flux adjustments, climatological surface fluxes were derived from the final 100 years of the spin-up run.

### **2.1.2 Atmosphere model**

The atmosphere model was initialised from a previous spin-up run, and was then integrated for 50 years under pre-industrial boundary conditions. The atmospheric carbon dioxide concentration was set to 280 ppm, the solar constant was set to  $1365 \text{ Wm}^{-2}$ , and modern (AD 1950, i.e. 0 years Before Present) values were used for the Earth's orbital parameters.

The bottom boundary condition was derived from the NOAA Optimum Interpolation v2 sea surface temperature analysis (*Reynolds et al.*, 2002), with climatological sea surface temperatures being calculated for the period 1982–2001. The ocean currents required by the sea ice model were diagnosed from the final 100 years of the ocean model spin-up run.

For the purposes of deriving flux adjustments, climatological surface fluxes were derived from the final 40 years of the spin-up run.

### **2.1.3 Coupled model**

The atmospheric and oceanic components of the model were initialised from their states at the end of the respective spin-up runs. The coupled model was then integrated for 100 years under pre-industrial boundary conditions, in order to overcome any initial coupling shock.

The mid-Holocene simulation was initialised from the state of the pre-industrial simulation at the end of year 100. Both simulations were then integrated for a further 100 years before beginning the PMIP2 experiments.

## **2.2 Experimental design**

The experimental design for the pre-industrial and mid-Holocene experiments is summarised in Table 1, and is discussed below.

Boundary condition	Pre-industrial	Mid-Holocene
Vegetation	Modern	Modern
Ice sheets	Modern	Modern
Topography/coastlines	Modern	Modern
CO <sub>2</sub> concentration [ppm]	280	280
CH <sub>4</sub> concentration [ppb]	-	-
N <sub>2</sub> O concentration [ppb]	-	-
Chlorofluorocarbons	-	-
O <sub>3</sub> concentration	Modern	Modern
Solar constant [Wm <sup>-2</sup> ]	1365	1365
Epoch [years BP]	0	6000
Eccentricity of Earth's orbit	0.016724	0.018682
Obliquity of Earth's axis [°]	23.446	24.105
Longitude of perihelion [°]	102.07	0.83

Table 1: Experimental design for the pre-industrial and mid-Holocene experiments.

### 2.2.1 Topography and bathymetry

Modern topography and bathymetry were used for both experiments.

The following straits are open on the ocean model grid, with the sill depths being as follows:

Greenland-Scotland Sill	800 m
Indonesian Throughflow	1010 m
Torres Strait	50 m
Bass Strait	50 m
Drake Passage	2800 m

The following straits are also open on the ocean model grid, with the sill depths being as follows; however, while water properties can be exchanged across these straits, *no net throughflow is allowed*:

Northwest Passage	50 m
Bering Strait	50 m
Hudson Strait	80 m
Strait of Gibraltar	270 m
Korea Strait	50 m
Mozambique Channel	1010 m

### 2.2.2 Land surface

The land surface model used modern distributions of vegetation and soil types for both experiments, as well as the modern positions of the ice sheets. The vegetation, soil types and ice sheets remained fixed throughout the experiments.

### 2.2.3 Radiative gases

An atmospheric CO<sub>2</sub> concentration of 280 ppm was used for both the pre-industrial and mid-Holocene experiments.

Modern ozone concentrations were also used for both experiments.

The radiation scheme in Mk3L does not account for the radiative effects of CH<sub>4</sub>, N<sub>2</sub>O or chlorofluorocarbons.

### 2.2.4 Insolation

The solar constant was set to 1365 Wm<sup>-2</sup> for both experiments.

Mk3L calculates the values of the Earth's orbital parameters at runtime, with the epoch being specified via the model's configuration file. The values calculated by the model for 0 and 6,000 years BP are shown in Table 1. The eccentricity and the obliquity of the Earth's axis agree, to the precision shown, with the values required by PMIP2 experimental design. The longitude of the perihelion differs from the values specified by PMIP2 by just +0.03° and -0.04° for the pre-industrial and mid-Holocene experiments respectively; these discrepancies are not considered to be significant.

## References

- Gordon, H. B., and S. P. O'Farrell (1997), Transient climate change in the CSIRO coupled model with dynamic sea ice, *Monthly Weather Review*, 125(5), 875–907.
- Gordon, H. B., L. D. Rotstayn, J. L. McGregor, M. R. Dix, E. A. Kowalczyk, S. P. O'Farrell, L. J. Waterman, A. C. Hirst, S. G. Wilson, M. A. Collier, I. G. Watterson, and T. I. Elliott (2002), The CSIRO Mk3 climate system model, *Technical Paper No. 60*, CSIRO Atmospheric Research, Aspendale, Victoria, Australia.
- Hirst, A. C., S. P. O'Farrell, and H. B. Gordon (2000), Comparison of a coupled ocean-atmosphere model with and without oceanic eddy-induced advection. Part I: Ocean spinup and control integration, *Journal of Climate*, 13, 139–163.
- Kanamitsu, M., W. Ebisuzaki, J. Woollen, S.-K. Yang, J. J. Hnilo, M. Fiorino, and G. L. Potter (2002), NCEP-DOE AMIP-II Reanalysis (R-2), *Bulletin of the American Meteorological Society*, 83(11), 1631–1643.
- National Oceanographic Data Center (2002), World Ocean Atlas 1998, URL [http://www.nodc.noaa.gov/OC5/pr\\_woa.html](http://www.nodc.noaa.gov/OC5/pr_woa.html).
- Phipps, S. J. (2006), The CSIRO Mk3L climate system model, *Technical Report No. 3*, Antarctic Climate & Ecosystems Cooperative Research Centre, Hobart, Tasmania, Australia, 236pp., ISBN 1-921197-03-X.

Reynolds, R. W., N. A. Rayner, T. M. Smith, D. C. Stokes, and W. Wang (2002), An improved in situ and satellite SST analysis for climate, *Journal of Climate*, 15, 1609–1625.